



Soils Map



©2026 AgriData, Inc.

Soils data provided by USDA and NRCS.



State: **Iowa**
 County: **Butler**
 Location: **29-93N-17W**
 Township: **Coldwater**
 Acres: **131.36**
 Date: **7/16/2026**

Maps Provided By:

 CUSTOMIZED ONLINE MAPPING
 © AgriData, Inc. 2025 www.AgriDataInc.com



Area Symbol: IA023, Soil Area Version: 32

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	*i Corn Bu	*i Alfalfa Tons	*i Soybeans Bu	*i Bluegrass Tons	*i Tall Grasses Tons	*i Water-Holding Inch	CSR2**	CSR
399	Readlyn silt loam, 1 to 3 percent slopes	39.70	30.1%		> 6.5ft.	Somewhat poorly drained	lw	220.8	5.7	64.0	4.0	6.6	11.3	91	90
84	Clyde silty clay loam, 0 to 3 percent slopes	31.29	23.8%		> 6.5ft.	Poorly drained	llw	224.0	4.7	65.0	4.0	6.7	11.7	88	75
198B	Floyd loam, 1 to 4 percent slopes	23.87	18.2%		> 6.5ft.	Somewhat poorly drained	llw	222.4	5.8	64.5	4.0	6.7	10.8	89	75
83B	Kenyon loam, 2 to 5 percent slopes	14.55	11.1%		> 6.5ft.	Moderately well drained	lle	225.6	6.3	65.4	4.1	6.8	10.1	90	85
175B	Dickinson fine sandy loam, 2 to 5 percent slopes	9.32	7.1%		> 6.5ft.	Well drained	llle	80.0	2.2	23.2	1.4	2.4	5.3	50	55

Soils data provided by USDA and NRCS.

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	*i Corn Bu	*i Alfalfa Tons	*i Soybeans Bu	*i Bluegrass Tons	*i Tall Grasses Tons	*i Water-Holding Inch	CSR2**	CSR
241C2	Saude-Burkhardt complex, 2 to 9 percent slopes, moderately eroded	5.15	3.9%		> 6.5ft.	Well drained	IVe	80.0	2.2	23.2	1.4	2.4	5.2	38	33
174B	Bolan loam, 2 to 5 percent slopes	2.57	2.0%		> 6.5ft.	Well drained	IIIs	177.6	5.0	51.5	3.2	5.3	8.6	64	70
782B	Donnan loam, 2 to 5 percent slopes	2.45	1.9%		> 6.5ft.	Somewhat poorly drained	IIe	80.0	2.1	23.2	1.4	2.4	10.0	43	50
391B	Clyde-Floyd complex, 1 to 4 percent slopes	1.70	1.3%		> 6.5ft.	Poorly drained	IIw	216.0	4.5	62.6	3.9	6.5	11.2	87	72
177	Saude loam, 0 to 2 percent slopes	0.76	0.6%		2.3ft. (Strongly contrasting textural stratification)	Well drained	IIIs	80.0	2.2	23.2	1.4	2.4	6.6	60	63
Weighted Average							1.85	202.5	5	58.7	3.6	6.1	10.4	83.2	76.9

**IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*c: Using Capabilities Class Dominant Condition Aggregation Method

	Symbol	Name	Description
	SAN	Sandy spot	A spot where the surface layer is loamy fine sand or coarser in areas where the surface layer of the named soils in the surrounding map unit is very fine sandy loam or finer. Typically .25 to 2 acres.