



# Schwartz Soils Map



State: **Minnesota**  
 County: **Yellow Medicine**  
 Location: **20-113N-38W**  
 Township: **Echo**  
 Acres: **144.78**  
 Date: **5/8/2026**



Soils data provided by USDA and NRCS.

Area Symbol: MN173, Soil Area Version: 20

Code	Soil Description	Acres	Percent of field	PI Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	Productivity Index	*n NCCPI Overall
421B	Amiret loam, 2 to 6 percent slopes	82.53	57.0%		> 6.5ft.	Well drained	Ile	98	83
954B2	Amiret-Swanlake loams, 2 to 6 percent slopes	29.94	20.7%		> 6.5ft.	Well drained	Ile	92	81
113	Webster clay loam, 0 to 2 percent slopes	16.89	11.7%		> 6.5ft.	Poorly drained	Ilw	93	83
L78A	Canisteo clay loam, 0 to 2 percent slopes	10.39	7.2%		> 6.5ft.	Poorly drained	Ilw	93	81
423	Seaforth loam, 1 to 3 percent slopes	3.38	2.3%		> 6.5ft.	Moderately well drained	Ils	95	84
134	Okoboji silty clay loam, 0 to 1 percent slopes	1.65	1.1%		> 6.5ft.	Very poorly drained	Illw	86	77
<b>Weighted Average</b>							<b>2.01</b>	<b>95.6</b>	<b>*n 82.4</b>

\*n: The aggregation method is "Weighted Average using all components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Symbol	Name	Description
WET	Wet spot	A somewhat poorly drained to very poorly drained area that is at least two drainage classes wetter than the named soils in the surrounding map unit. Typically 0.5 to 2 acres.

Soils data provided by USDA and NRCS.