



Soils Map



©2026 AgriData, Inc.

Soils data provided by USDA and NRCS.



State: **Iowa**
 County: **Monroe**
 Location: **26-73N-19W**
 Township: **Cedar**
 Acres: **28.75**
 Date: **6/12/2026**



Maps Provided By:



© AgriData, Inc. 2025

www.AgriDataInc.com



Area Symbol: IA135, Soil Area Version: 33

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	*i Corn Bu	*i Alfalfa Tons	*i Soybeans Bu	*i Bluegrass Tons	*i Tall Grasses Tons	*i Water-Holding Inch	CSR2**	CSR	*n NCC Over
423D2	Bucknell silty clay loam, 9 to 14 percent slopes, moderately eroded	15.43	53.6%		0.7ft. (Abrupt textural change)	Somewhat poorly drained	IVe	97.6	2.5	28.3	1.8	2.9	9.4	6	13	
731C2	Pershing silty clay loam, 5 to 9 percent slopes, moderately eroded	10.23	35.6%		> 6.5ft.	Somewhat poorly drained	IIle	80.0	2.1	23.2	1.4	2.4	11.8	62	45	
131B	Pershing silt loam, 2 to 5 percent slopes	2.78	9.7%		> 6.5ft.	Somewhat poorly drained	IIle	80.0	2.1	23.2	1.4	2.4	11.8	70	67	
13B	Olmitz-Colo-Vesser complex, 2 to 5 percent slopes	0.31	1.1%		> 6.5ft.	Poorly drained	IIw	192.0	4.0	55.7	3.5	5.8	11.7	82	60	
Weighted Average								3.53	90.7	2.3	26.3	1.6	2.7	10.5	30.1	*n

Soils data provided by USDA and NRCS.



**IA has updated the CSR values for each county to CSR2.

*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

	Symbol	Name	Description
	CLR	Clay spot, red	An exposure of clay that formed in red truncated paleosol weathered from glacial till, surrounded by soils that are not clayey. Textures included would be clay and silty clay with clay contents greater than 40 percent. Typically .25 to 2 acres.