



Soils Map



©2026 AgriData, Inc.

Soils data provided by USDA and NRCS.



State: **Iowa**
 County: **Wapello**
 Location: **22-72N-14W**
 Township: **Center**
 Acres: **59.42**
 Date: **7/6/2026**



Maps Provided By:



© AgriData, Inc. 2025

www.AgriDataInc.com



Area Symbol: IA179, Soil Area Version: 34

Code	Soil Description	Acres	Percent of field	CSR2 Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	*j Corn Bu	*j Alfalfa Tons	*j Soybeans Bu	*j Bluegrass Tons	*j Tall Grasses Tons	*j Water-Holding Inch	CSR2**	CSR	*N C
831C2	Pershing silty clay loam, terrace, 5 to 9 percent slopes, moderately eroded	16.76	28.3%		> 6.5ft.	Somewhat poorly drained	Ille	80.0	2.1	23.2	1.4	2.4	11.8	64	40	
S732C2	Weller silty clay loam, 5 to 9 percent slopes, moderately eroded	15.23	25.6%		> 6.5ft.	Moderately well drained	Ille	0.0	0.0	0.0	0.0	0.0	0.0	59		
132C	Weller silt loam, 5 to 9 percent slopes	11.35	19.1%		> 6.5ft.	Moderately well drained	Ille	80.0	2.2	23.2	1.4	2.4	10.6	59	35	
832C2	Weller silty clay loam, terrace, 5 to 9 percent slopes, moderately eroded	5.72	9.6%		> 6.5ft.	Moderately well drained	Ive	80.0	2.2	23.2	1.4	2.4	10.6	61	30	

Soils data provided by USDA and NRCS.



Code	Soil Description	Acres	Percent of field	CSR2 Legend	Restrictive Layer	Soil Drainage	Non-Irr Class *c	*i Corn Bu	*j Alfalfa Tons	*i Soybeans Bu	*j Bluegrass Tons	*i Tall Grasses Tons	*j Water-Holding Inch	CSR2**	CSR	*i N C	
880C2	Clinton silt loam, terrace, 5 to 9 percent slopes, eroded	3.23	5.4%		> 6.5ft.	Moderately well drained	IIIe	187.2	5.2	54.3	3.4	5.6	11.5	71	60		
65F	Lindley loam, 18 to 25 percent slopes	3.00	5.0%		> 6.5ft.	Well drained	VIIe	115.2	3.2	33.4	2.1	3.5	9.4	17	10		
425D2	Keswick loam, 9 to 14 percent slopes, moderately eroded	1.89	3.2%		0.8ft. (Abrupt textural change)	Somewhat poorly drained	IVe	88.0	2.5	25.5	1.6	2.6	8.8	9	5		
795D2	Ashgrove silty clay loam, 9 to 14 percent slopes, moderately eroded	1.06	1.8%		> 6.5ft.	Poorly drained	IVe	88.0	1.8	25.5	1.6	2.6	8.4	7	7		
W	Water	0.75	1.3%		> 6.5ft.			0.0	0.0	0.0	0.0	0.0	0.0	0	0		
594D2	Galland loam, heavy loess, 9 to 14 percent slopes, moderately eroded	0.35	0.6%		> 6.5ft.	Somewhat poorly drained	IVe	88.0	2.3	25.5	1.6	2.6	9.7	19	5		
730B	Nodaway-Cantril complex, 2 to 5 percent slopes	0.08	0.1%		> 6.5ft.	Moderately well drained	IIW	209.6	5.4	60.8	3.8	6.3	11.4	80	63		
Weighted Average								*-	66.7	1.8	19.3	1.2	2	8	55.7	*-	

**IA has updated the CSR values for each county to CSR2.

*- CSR weighted average cannot be calculated on the current soils data, use prior data version for csr values.*i Yield data provided by the ISPAID Database version 8.1.1 developed by IA State University.

*n: The aggregation method is "Weighted Average using all components"

*c: Using Capabilities Class Dominant Condition Aggregation Method

*- Non Irr Class weighted average cannot be calculated on the current soils data due to missing data.