


B895 SC Newberry County
188.08 Acres



B895 SC Newberry County

Soil Report

Symbol	Map Unit Description	Farmland class	Area (AC)	Available Water Storage (0-150cm - Weighted Average)	Flooding Frequency (Maximum)	Drainage Class (Dominant Condition)	Drainage Class (Wettest Component)	Hydric Classification	Water Table Depth (Annual Minimum)	Water Table Depth (April to June Minimum)	Bedrock Depth (Minimum)	Soil Order
SeB	Sedgefield sandy loam, 0 to 4 percent slopes	Farmland of statewide importance	0	24.0	None	Somewhat poorly drained	Somewhat poorly drained	0	33	0	0	Alfisols
WnB	Winnboro sandy loam, 2 to 6 percent slopes	Farmland of statewide importance	0	12.0	None	Well drained	Well drained	0	0	0	104	Alfisols
Wyc2	Wynott-Winnboro complex, 6 to 10 percent slopes, moderately eroded	Not prime farmland	0	10.0	None	Well drained	Well drained	0	0	0	74	Alfisols
HaB	Hard Labor sandy loam, 2 to 6 percent slopes	All areas are prime farmland	0	18.3	None	Moderately well drained	Moderately well drained	0	114	0	0	Ultisols
CnA	Chenneby silt loam, 0 to 2 percent slopes, frequently flooded	Prime farmland if drained and either protected from flooding or not frequently f	0	27.3	Frequent	Somewhat poorly drained	Somewhat poorly drained	5	51	0	0	Inceptisols
SaB	Santuc loamy coarse sand, 2 to 6 percent slopes	All areas are prime farmland	0	22.3	None	Moderately well drained	Moderately well drained	0	66	0	0	Ultisols