


B935 - SC Laurens
22.4 Acres



B935 - SC Laurens

Soil Report

| Symbol | Map Unit Description | Farmland class | Area (AC) | Available Water Storage (0-150cm - Weighted Average) | Flooding Frequency (Maximum) | Drainage Class (Dominant Condition) | Drainage Class (Wettest Component) | Hydric Classification | Water Table Depth (Annual Minimum) | Water Table Depth (April to June Minimum) | Bedrock Depth (Minimum) | Soil Order |
|--------|--|--|-----------|--|------------------------------|-------------------------------------|------------------------------------|-----------------------|------------------------------------|---|-------------------------|------------|
| PaD2 | Pacolet sandy clay loam, 10 to 15 percent slopes, eroded | Not prime farmland | 0 | 19.1 | None | Well drained | Well drained | 0 | 0 | 0 | 0 | Ultisols |
| Ca | Cartecay-Toccoa complex | Prime farmland if drained and either protected from flooding or not frequently f | 0 | 14.7 | Occasional | Somewhat poorly drained | Somewhat poorly drained | 5 | 31 | 31 | 0 | Entisols |
| CeB2 | Cataula sandy clay loam, 2 to 6 percent slopes, eroded | Farmland of statewide importance | 0 | 19.0 | None | Moderately well drained | Moderately well drained | 0 | 94 | 94 | 0 | Ultisols |
| CeC2 | Cataula sandy clay loam, 6 to 10 percent slopes, eroded | Not prime farmland | 0 | 19.0 | None | Moderately well drained | Moderately well drained | 0 | 94 | 94 | 0 | Ultisols |
| EnD | Enon sandy loam, 10 to 15 percent slopes | Not prime farmland | 0 | 19.8 | None | Well drained | Well drained | 0 | 0 | 0 | 0 | Alfisols |
| CdB2 | Cataula sandy loam, 2 to 6 percent slopes, eroded | Farmland of statewide importance | 0 | 18.5 | None | Moderately well drained | Moderately well drained | 0 | 94 | 94 | 0 | Ultisols |
| WIF | Wilkes soils, 15 to 40 percent slopes | Not prime farmland | 0 | 3.6 | None | Well drained | Well drained | 0 | 0 | 0 | 25 | Alfisols |